



HIV AMONG WHITES IN MARYLAND, 2012

OCTOBER 2014

New HIV Diagnoses

- Non-Hispanic (NH) whites accounted for 13.2% (193) of the 1,466 reported adult/adolescent HIV diagnoses (age 13+) in Maryland in 2012.
- The proportion of HIV diagnoses in Maryland that were NH white has decreased from 38.3% in 1985 to 13.2% in 2012.
- Case rates among newly reported diagnoses in 2012 remain lowest among NH whites (7.0 per 100,000) as compared to NH blacks (79.3) and Hispanics (22.4).





Characteristics of Whites Living with HIV

Age Categories

- 32.3% 40's
- 31.5% 50's
- 15.3% 30's
- 14.4% 60+
- 5.9% 20's
- 0.4% Teens

İ

Exposure Category

- 56.9% MSM
- 20.4% IDU
- 18.1% HET
- Of the 29,553 total living HIV cases (with or without AIDS) on 12/31/2012 in Maryland, 14.9% (4,399) of those were among NH whites. In other words, 1 in 623 NH whites were reported to be living with HIV in Maryland.
- Total living HIV case rates in Maryland were 3 times higher among NH white males (218.7 cases per 100,000 on 12/31/2012, all ages) as compared to NH white females (65.0). In other words, 1 in 457 NH white males and 1 in 1,538 NH white females in Maryland were reported to be living with HIV in Maryland.
- Total living HIV case rates among NH whites in Maryland were highest among males in their 40's (473.2 cases per 100,000 on 12/31/2012, all ages) followed by men in their 50's (452.6).
- Close to three quarters (72.3%) of the NH white men diagnosed to be living with HIV in Maryland were among Men who have Sex with Men (MSM). Among NH white women reported to be living with HIV in Maryland,
 54.3% were attributed to Heterosexual Exposure (HET) followed by Injection Drug Users (IDU) (44.6%).

MSM: Men who have Sex with Men, HET: Heterosexual Exposure; IDU: Injection Drug User